

Table 10.1
Kaposi Sarcoma

Number of Cases

By SEER Location and Year of Diagnosis, 1975-2006

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2006
SEER 9 ^b	134	79	326	887	1573	2131	2233	2056	1801	838	390	398	329	503
SEER 9 except SF-Oakland	102	44	85	259	427	775	842	975	874	462	243	264	221	347
San Francisco-Oakland SMSA ^c	32	35	241	628	1146	1356	1391	1081	927	376	147	134	108	156
San Francisco City/County	15	21	201	540	955	1077	1089	792	690	253	86	79	68	103

Age-Adjusted Cancer Incidence Rates^a

By SEER Location and Year of Diagnosis, 1975-2006

All Races, Males, All Ages

Year of Diagnosis

SEER Location	1975-1979	1980-1981	1982-1983	1984-1985	1986-1987	1988-1989	1990-1991	1992-1993	1994-1995	1996-1997	1998-1999	2000-2001	2002-2003	2004-2006
SEER 9 ^b	0.4	0.5	1.7	4.2	7.1	9.2	9.3	8.2	7.0	3.2	1.5	1.5	1.2	1.2
SEER 9 except SF-Oakland	0.4	0.4	0.6	1.5	2.3	4.0	4.2	4.6	4.0	2.1	1.1	1.2	1.0	1.0
San Francisco-Oakland SMSA ^c	0.7	1.3	7.0	17.2	30.0	34.0	34.0	25.6	22.0	8.9	3.5	3.0	2.6	2.4
San Francisco City/County	1.1	3.1	25.1	65.3	111.3	124.6	125.7	89.1	77.5	28.2	9.4	8.5	7.4	7.2

^a Rates are per 100,000 males and are age-adjusted to the 2000 US Std Population (19 age groups - Census P25-1130).^b The SEER 9 areas are San Francisco-Oakland SMSA, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah and Atlanta.^c The San Francisco-Oakland SMSA represents 15% of all persons in the SEER areas. For this age group, all SEER areas may not be generalizable to the United States for Kaposi Sarcoma.

- Statistic not shown. Rate based on less than 4 cases for the time interval.